

Research



Cite this article: Aleta A, Hisi ANS, Meloni S, Poletto C, Colizza V, Moreno Y. 2017 Human mobility networks and persistence of rapidly mutating pathogens. *R. Soc. open sci.* **4:** 160914.
<http://dx.doi.org/10.1098/rsos.160914>

Received: 16 November 2016

Accepted: 10 February 2017

Subject Category:

Biology (whole organism)

Subject Areas:

mathematical modelling/theoretical biology/computer modelling and simulation

Keywords:

pathogen persistence, spatial epidemic spread, metapopulation model, susceptible-infected-recovered-susceptible, waning of immunity

Author for correspondence:

Chiara Poletto
e-mail: chiara.poletto@inserm.fr

[†]These authors contributed equally to this study.

[‡]Present address: 91, Avenue du Général Leclerc 92340, Bourg la Reine, France

Human mobility networks and persistence of rapidly mutating pathogens

Alberto Aleta^{1,†}, Andreia N. S. Hisi^{3,†,‡}, Sandro Meloni^{1,2}, Chiara Poletto³, Vittoria Colizza^{3,4} and Yamir Moreno^{1,2,4}

¹Institute for Biocomputation and Physics of Complex Systems, and ²Department of Theoretical Physics, University of Zaragoza, Zaragoza, Spain

³Sorbonne Universités, UPMC Univ Paris 06, INSERM, Institut Pierre Louis d'Épidémiologie et de Santé Publique (IPLESP UMRS 1136), Paris, France

⁴ISI Foundation, Turin, Italy

SM, 0000-0001-6202-3302; CP, 0000-0002-4051-1716

Rapidly mutating pathogens may be able to persist in the population and reach an endemic equilibrium by escaping hosts' acquired immunity. For such diseases, multiple biological, environmental and population-level mechanisms determine the dynamics of the outbreak, including pathogen's epidemiological traits (e.g. transmissibility, infectious period and duration of immunity), seasonality, interaction with other circulating strains and hosts' mixing and spatial fragmentation. Here, we study a susceptible-infected-recovered-susceptible model on a metapopulation where individuals are distributed in sub-populations connected via a network of mobility flows. Through extensive numerical simulations, we explore the phase space of pathogen's persistence and map the dynamical regimes of the pathogen following emergence. Our results show that spatial fragmentation and mobility play a key role in the persistence of the disease whose maximum is reached at intermediate mobility values. We describe the occurrence of different phenomena including local extinction and emergence of epidemic waves, and assess the conditions for large-scale spreading. Findings are highlighted in reference to previous studies and to real scenarios. Our work uncovers the crucial role of hosts' mobility on the ecological dynamics of rapidly mutating pathogens, opening the path for further studies on disease ecology in the presence of a complex and heterogeneous environment.

1. Introduction

Many pathogens are able to persist in a host population by repeatedly reinfecting individuals who lose the immunity

acquired during infection [1]. This mechanism is particularly important for the case of rapidly mutating pathogens when mutation is associated with antigenic drift—e.g. for influenza virus [2] and respiratory syncytial virus [3]. The probability of pathogen persistence is then determined by the interplay between the transmissibility and infection time scales (i.e. duration of infection and immunity period), along with ecological factors like seasonality, interaction with other circulating pathogens, mixing patterns among individuals, their spatial distribution and mobility.

Waning of immunity and partial immunization have been accounted for by several modelling studies [4–9] addressing detailed immunological mechanisms [10], interaction with other strains [7, 11], human contact structure [6], environmental factors [12], seasonality and temporal variation in transmission [13–15]. The fate of an emerging pathogen in terms of persistence/extinction was also studied, both as a theoretical problem [8,9] and through applications to the emergence of a novel influenza strain [2,16]. However, the vast majority of studies on waning of immunity and persistence have disregarded spatial structure so far.

Recently, it has been shown that the geographical distribution of the host population and the rate of travelling among distant locations determine the time scale of human mixing, hence affecting the probability of having large-scale epidemic spread [17–20] and the incidence profile [21–24]. For the case of childhood diseases (e.g. measles), previous studies showed that these dynamical effects may impact disease persistence and alter the critical community size [25–31], i.e. the population threshold above which a disease is able to persist [25]. Results reported so far, however, suggest that the persistence probability is determined by the interplay between the different spatial ingredients—e.g. distance and frequency of travels, environmental influences on transmission—in a non-trivial way. Indeed, different contrasting hypotheses have been suggested, with persistence maximized for intermediate levels of spatial coupling [28,30], for maximum levels of coupling [31] or either of the two depending of the level of spatial heterogeneity in transmission [29].

The spatial structure of human population could have important and non-trivial effects on the spreading of diseases also when waning of immunity is at work, affecting the possible dynamical regimes following an outbreak. In this paper, we address this problem by means of a stochastic discrete spatially explicit metapopulation model. We reconstruct the dynamics of an emerging infection through extensive numerical simulations. We characterize the role played by the spatial fragmentation, the coupling, and the structure of the mobility network through which individuals travel from one sub-population to another. We focus on the regime of parameters that best describes influenza-like infections. Our metapopulation framework realistically reproduces the statistical properties of the human population, i.e. strong heterogeneities and correlations in the population distribution and mobility. We perform a detailed analysis of the mechanisms shaping the probability of pathogen persistence and the spatial dynamics following emergence in the short and long term. Our approach offers the theoretical and computational framework to further explore the effects of other features not taken into account here (e.g. multiple strains, seasonality and others) in order to gain a full understanding of pathogens' persistence and be able to perform realistic data-driven simulations.

2. Material and methods

We use a stochastic individual-based metapopulation model [32–34], where mobility and infection of individuals are explicitly accounted for as discrete-time processes. Individuals mix homogeneously within the local communities (also called sub-populations, patches or nodes of the metapopulation system), whereas at the global level the coupling between these sub-populations is introduced by individuals that travel along mobility connections (figure 1). The model is thus represented in terms of a network of links, i.e. mobility flows, connecting different nodes representing local populations.

Massive data and extensive research studies have uncovered the statistical properties of the distribution of human populations and their associated mobility, pointing out their network structure, characterized by large variabilities and non-trivial correlations [35–38]. In particular, the population of urban areas and the number of connections between locations (considering e.g. air-transportation or commuting) span several orders of magnitude, and more-populated centres are also often the hubs of the mobility network. We account for these properties by assuming a power law distribution for the patches' connectivity and a linear relationship between connectivity and population. Specifically, we considered a network of V patches, each of them (node i of the network) characterized by a population of size N_i , and a degree k_i that represents the number of sub-populations through which node i is linked via mobility connections. The nodes' degree distribution is assumed to be of the form $P(k) \sim k^{-\gamma}$ with




Figure 1. Schematic representation of the stochastic metapopulation model. Panel (a) represents the disease dynamics that follows a SIRS scheme, where individuals change states with given transition probabilities (Susceptible (S), Infected (I), Recovered (R) and back to S when the acquired immunity is lost). Panel (b) illustrates the structure of the metapopulation network, which consists of sub-populations that are connected by individuals travelling from one to another through a mobility network.

no degree–degree correlation present; a random network with these characteristics was generated by means of the uncorrelated configuration model [39] with $\gamma = 3.0$. Mobility fluxes are assigned to each link by assuming that, in each sub-population, individuals may travel at each time step to neighbouring sub-populations with a probability p . Departing individuals from node i choose at random one of the available k_i links, so that the probability of travelling along a connection is given by p/k_i . This process yields a population distribution that at equilibrium is proportional to the number of connections, namely, $N_i = k_i \langle N \rangle / \langle k \rangle$; where $\langle N \rangle$ and $\langle k \rangle$ are the average population per node and average degree of the metapopulation network, respectively [20].

The infection dynamics is implemented through the Susceptible-Infected-Recovered-Susceptible (SIRS) compartmental scheme [4]. The evolution of the dynamics results from the following transition rules iterated at each time step, corresponding to 1 day:

- (i) Susceptible individuals within each sub-population i may be infected with probability $\Gamma_i = 1 - (1 - \beta/N_i)^{I_i}$, where β represents the transmission probability, and N_i and I_i denote, respectively, the total population and the number of infected individuals in node i ;
- (ii) Infected individuals enter with probability μ the recovered compartment, that represents a state in which individuals are immune to the disease; here immunity is assumed to be complete;
- (iii) Eventually, recovered individuals lose their immunity and get back to the susceptible state. This occurs with probability λ , that is the inverse of the average immunity period, L .

Infection transition rules combined with travelling can be summarized, for each sub-population, by the following discrete master equations:

$$S_i(t + dt) - S_i(t) = (1 - p)[\lambda R_i(t) - \Gamma_i(t)S_i(t)] - pS_i(t) + \sum_{j \in v_i} \frac{p}{k_j} [S_j(t) + \lambda R_j(t) - \Gamma_j(t)S_j(t)]$$

$$I_i(t + dt) - I_i(t) = (1 - p)[\Gamma_i(t)S_i(t) - \mu I_i(t)] - pI_i(t) + \sum_{j \in v_i} \frac{p}{k_j} [I_j(t)(1 - \mu) + \Gamma_j(t)S_j(t)]$$

$$R_i(t + dt) - R_i(t) = (1 - p)[\mu I_i(t) - \lambda R_i(t)] - pR_i(t) + \sum_{j \in v_i} \frac{p}{k_j} [R_j(t)(1 - \lambda) + \mu I_j(t)],$$

where v_i represents the set of neighbouring patches of i . The described process disregards vital dynamics. This choice simplifies the dynamical characterization of the model and it is motivated by our interest in diseases with durations of both infection and immunity periods much shorter than the average life duration (e.g. influenza). Starting with a fully susceptible population, the disease is seeded in a randomly chosen node by infecting 0.5% of its population, in such a way that stochastic runs differ for both the initial seed and the stochastic realization of the dynamics. The disease may then propagate inside the local population and spread to neighbouring populations due to host movements. Progression of the disease and host movements are stochastic events that are numerically modelled through multinomial processes. For each scenario considered, 10^3 independent epidemics are simulated starting from different seeded sub-populations.

We consider influenza-like infections spreading in space as a prototypical example for our study. We fix the duration of infection $\mu^{-1} = 2.5$ days and the basic reproductive number $R_0 = \beta/\mu = 2.0$. The population is assumed to be composed of 10^8 individuals divided in V patches. We explored scenarios characterized by: (i) different spatial fragmentation of the host population, considering $V = 10^2, 10^3, 10^4$; (ii) a varying coupling between patches expressed by host mobility, considering $p \in [10^{-6}, 1]$; (iii) different topological structures of mobility connections, comparing the heterogeneous network described by a power-law degree distribution to a homogeneous Erdős–Rényi graph [40] described by a Poisson degree distribution with the same average value; (iv) a varying immunity period, considering $L \in [1, 800]$ days.

For each set of parameters, we numerically monitor the global prevalence (i.e. the total fraction of infectious individuals) and the number of infected patches $D(t)$ in time. We also estimate the probability that the pathogen persists in the population at the global level, P_{gl} , by computing the fraction of stochastic runs for which the epidemic reaches the endemic state, once a given set of parameter values is considered. The same quantity can be defined at the local level to measure the probability of local persistence P_{loc} . This can be computed considering a closed population, or a population integrated into a metapopulation framework, thus open to incoming and outgoing fluxes of individuals. At the local level, we also define an outbreak as a single epidemic (infecting at least 0.1% of the population) that continuously persists in the population. Repeated outbreaks instead are considered as distinct waves following local extinction. These behaviours are characterized for the different scenarios explored, in order to provide a coherent picture of the mechanisms for virus persistence and endemic equilibrium in the spatially structured population.

3. Results and discussion

In the following, we present first the results regarding the persistence of the pathogen in the population and the mechanisms behind it, followed by a detailed characterization of the spatial dynamics across the scenarios explored.

3.1. Persistence

3.1.1. Interplay between waning of immunity and mobility

The probability of persistence is found to be very high and equal to 1 for short immunity periods; once immunity becomes longer, a reduction of the probability of persistence is observed (figure 2a). This behaviour is common to all host mobility rates explored. When the waning of immunity is very fast, there is a high rate of renewal of susceptible individuals that can be newly infected by the pathogen, thus ensuring its survival in the population. A slower rate of renewal makes persistence less likely and the specific behaviour of the decrease in persistence probability depends on the coupling between patches. Reduction may occur through a drop to zero persistence (e.g. $p = 10^{-6}, 10^{-5}$, etc. and $p = 10^{-2}, 10^{-1}$, etc.), or a drop to an intermediate positive value of persistence probability that is maintained for a large range of immunity period, before reaching $P_{\text{gl}} = 0$ (e.g. $p = 10^{-4}, 2 \times 10^{-4}$, etc.).

The dependence of persistence drop on mobility is highlighted in figure 2b. Our results show that it is possible to distinguish three mobility regimes—namely, low mobility (cyan), intermediate mobility (light blue) and high mobility (purple). These regimes are approximately delimited by $p < 4.5 \times 10^{-5}$, $p \in [4.5 \times 10^{-5}, 2.0 \times 10^{-3}]$ and $p > 2.0 \times 10^{-3}$ (referring to the profile of 5% persistence probability), respectively.




Figure 2. (a) Persistence probability P_{gi} as a function of the duration of immunity L for several values of the travelling probability p . Three different regions can be distinguished: low mobility $10^{-6} \leq p \leq 4.5 \times 10^{-5}$ (cyan), intermediate values $4.5 \times 10^{-5} \leq p \leq 10^{-3}$ (light blue) and high mobility $2 \times 10^{-3} \leq p \leq 1$ (purple). (b) Colour map of P_{gi} in the (p, L) plane. Black curves highlight the contours with $P_{gi} = 5\%$ and $P_{gi} = 95\%$ as indicated in the figure. The red arrow indicates the value of the global invasion threshold for the mobility network and disease parameters here considered, $\mu R_0^2 / [2 \langle N \rangle (R_0 - 1)^2] \langle k \rangle^2 / (\langle k^2 \rangle - \langle k \rangle) = 4.5 \times 10^{-5}$. Both panels show results obtained with $V = 10^4$, $\langle N \rangle = 10^4$ and power law degree distribution.

In the low mobility regime, persistence at the global level is mainly determined by the probability that the pathogen would persist locally in the seeded population. If p is low enough so that no infectious individual travels out of the infected patch during the first wave of the epidemic, pathogen survival following the first wave depends on the length of the immunity period. A long enough immunity period prevents the replenishment of healthy individuals in the short term, thus leading to the extinction of the epidemic in the seeded patch, with no chances for global spread. If the availability of new susceptible hosts is instead provided fast enough by a short immunity period, the epidemic can survive the first wave in the seeded population, reaching a local endemic equilibrium. This, in its turn, can slowly seed the epidemic in neighbouring patches even when small mobility rates are considered, thanks to the local equilibrium condition. This same process is then occurring in each newly seeded patch, so that propagation unfolds at the global level and a high probability of persistence is observed.

The condition distinguishing between an epidemic trapped locally in a patch and one that spreads at the global level has been quantified in previous studies in terms of the global invasion threshold [17,20, 41–45]. Its analytical expression, provided in [20], provides an upper bound for the low mobility regime

$$p < \frac{1}{\langle N \rangle} \frac{\langle k \rangle^2}{\langle k^2 \rangle - \langle k \rangle} \frac{\mu R_0^2}{2(R_0 - 1)^2}. \quad (3.1)$$

Once informed with the demographic and topological features of our spatially structured population, this condition is found to provide a good approximation to the limit of the low mobility regime (red arrow in figure 2b).

The spatial structure of the population becomes less important once the system is found in the high mobility regime. In this case, the probability of moving from one patch to another is so high as to largely increasing the opportunity of mixing of the populations belonging to different patches. The system behaves effectively as a single population of the size of the full metapopulation (i.e. in our case 10^8 individuals) affected by a SIRS dynamics. The conditions for global persistence are therefore those described before for the seeded population in the low mobility regime. Differently from before, here local persistence is equivalent to global persistence, due to the large degree of mixing across space. The values of the immunity period at which a given persistence probability is observed (e.g. $L \simeq 300$ for $P_{gl} = 5\%$) differ from the case of low mobility regime ($L \simeq 100$), because of the difference in the population size, i.e. a single patch in the low mobility regime and effectively the full metapopulation size in the high mobility regime.

The intermediate mobility regime displays a higher persistence probability for a given duration of the immunity period, compared with low and high mobility regimes (figure 2b). The contour line at a fixed P_{gl} (e.g. $P_{gl} = 5\%$ or 95%) exhibits a maximum for intermediate values of the travelling probability p that is reached for high values of L . In addition, a smoother variation of the persistence probability is observed by varying L for a given p , compared with the faster transitions recovered in the low and high mobility regimes. The intermediate degree of mixing provided by intermediate mobility seems to enhance stochastic effects responsible for local extinction, thus having an important impact on the persistence at the global level.

Maximum persistence for intermediate level of coupling was also reported in the context of childhood diseases, where SIR models with continuous external introduction of the infection were considered [28,30]. No consensus was, however, found in this context, with other studies reporting different findings. For instance, Hagensaar *et al.* found an increase in the persistence with increasing levels of the coupling among sub-populations [31]. Rozhnova *et al.* confirmed this finding when homogeneous transmission across patches is considered, and additionally showed that heterogeneity in transmission among patches selects dynamical regimes with different spatial effects [29]. Different results obtained from different models show how the detailed mechanism of susceptible replenishment, together with infectious transmission and recovery, and spatial mixing, have an impact on the resulting persistence. Prompted by this, we assess in the following the role played by the spatial fragmentation of the population and by the mobility network on our model where the replenishment of individuals is obtained through immunity waning.

3.1.2. Role of spatial fragmentation and mobility network

Spatial fragmentation is explored by subdividing the total population of $N = 10^8$ hosts into $V = 10^2, 10^3, 10^4$ patches. Global persistence in the three scenarios shows the same qualitative behaviour described in the previous section (figure 3a). The separation between different mobility regimes, however, depends on the number of patches considered. More precisely, it is a demographic effect due to a varying average population size per patch. Referring to the transition from low mobility to intermediate mobility regime, we have indeed that the estimate expressed in equation (3.1) depends on $\langle N \rangle$. Increasing the number of patches from $V = 10^2$ to $V = 10^3, 10^4$, the average size of the population in a given patch decreases by one and two orders of magnitude, respectively, thus inducing a corresponding increase in the mobility value of the transition from low to intermediate regimes. As a result, the maximum of the global persistence is obtained for correspondingly increasing values of the travelling probability.

It is interesting to note that the behaviour observed in the high mobility regime is the same across the three scenarios. This confirms that the resulting dynamics when p is high enough corresponds effectively to the local persistence dynamics of a single population of 10^8 individuals, as discussed in the previous subsection.

If the topological structure connecting the patches via mobility links is altered and homogenized, we find that higher values of mobility need to be reached for the transition from low to intermediate mobility to occur (figure 3b). As expected, heterogeneous mobility patterns (as the ones adopted in figure 2) favour hosts' mobility resulting in a larger global persistence for lower travelling probabilities. This is consistent with what is predicted by the global invasion of equation (3.1) as shown by the arrow in figure 3b. At high mobility regimes, the structure connecting the links does not play any role, and global persistence is the




Figure 3. (a) Fragmentation analysis. Curves represent $P_{gl} = 5\%$ for three values for the number of patches $V = 10^2, 10^3$ and 10^4 sub-populations. In all the cases, the total population has been kept constant $N_{tot} = 10^8$ ending up with an average population per sub-population of $\langle N \rangle = 10^6, 10^5$ and 10^4 , respectively. Other parameters as in figure 2. (b) Impact of network topology. Curves represent $P_{gl} = 5\%$ for homogeneous (Erdős–Rényi graph [40], green line) and heterogeneous (power law distribution, blue line) mobility networks. Both networks have the same number of patches $V = 10^4$ and similar average degree $\langle k \rangle$. Other parameters as in figure 2.

same. Mixing is high enough so that the structured population behaves effectively as a single population regardless of the topology through which such mixing occurs.

3.2. Spatial dynamics

In this section, we go more in depth into the understanding of the spatial dynamics following the emergence of a pathogen in the fully susceptible and spatially structured population. Starting from the global persistence diagram reported in figure 2, we characterize the epidemic diffusion in space considering a single value of reference for p in each of the three mobility regimes ($p = 3 \times 10^{-5}$ in the low, $p = 6 \times 10^{-4}$ in the intermediate, and $p = 10^{-2}$ in the high mobility regime), associated with different values of the immunity period L (figure 4).

3.2.1. Low mobility coupling

We first analyse the low mobility regime, by characterizing the spatial epidemic dynamics obtained for $p = 3 \times 10^{-5}$ and $L = 30$ leading to $P_{gl} = 1.0$ (figure 4g), and $p = 3 \times 10^{-5}$ and $L = 80$ leading to $P_{gl} = 0.50$ (figure 4d). As discussed previously, global persistence is determined by local persistence of the disease at the patch level. The very high persistence observed for $p = 3 \times 10^{-5}$ and $L = 30$ is ensured by an endemic equilibrium that is reached in all patches after a given time, as signalled by 100% of patches experiencing an outbreak (i.e. $D(t)/V = 1$ after a given t). Even with a low probability of moving from one patch to another, the initially seeded population is able to transfer infected hosts to neighbouring populations in the long time limit thanks to the endemic equilibrium reached. In their turn, newly infected patches can




Figure 4. Spatial dynamics. The external panels show the time evolution of the fraction of infected sub-populations $D(t)/V$ for different values of the duration of immunity L and travelling probability p . The central panel shows the different regions where each value of L and p has been chosen. In each panel, curves corresponding to 10^2 runs with different initial conditions are shown. To facilitate visualization, 90% of them are smoothed.

further propagate to the infection reaching all patches of the system, thus achieving a high probability of persistence of the pathogen at the global level, fuelled by each endemic dynamics.

The same travelling probability leads to lower P_{gl} once the immunity period increases, from $L = 30$ to $L = 80$ (figure 4d). Here, a similar mechanism of local persistence and spatial diffusion ensured by the endemic equilibrium is at play, but it does not occur in all patches. At the end of our simulations, we find indeed that only a fraction of patches is experiencing an outbreak ($D/V \simeq 80\%$). The different dynamics occurring across patches depends on the patch population size. For a fixed value of the immunity period L , there exists a minimum population N_{min} needed in order to achieve a certain probability of local persistence (figure 5) [8,25,46,47]. The population size of the patch of our system varies approximately in the range $[5 \times 10^3, 2.7 \times 10^5]$, due to the heterogeneous topology connecting the patches. If $L = 30$, we find that all population values are above the minimum population needed to achieve local persistence and mobility can act as spatial spreader to reach global persistence. If $L = 80$, we find that a portion of the patches do not have enough population to achieve local persistence. In particular, to obtain a local persistence probability higher than 95% the minimum population size is 5×10^4 . The epidemic is able to spread at the global scale; however, smaller peripheral patches do not meet the condition for local endemic equilibrium. This results in a slower spatial spread, compared with the case of $L = 30$ (as displayed by the time axis in figure 4d,g), and an epidemic that is not able to endemically reach all sub-populations ($D/V < 1$).

The dynamics for the $L = 80$ case reflects a spatial hierarchy similar to the one studied in the context of childhood diseases—notably, for measles epidemics in the UK [23,24,48]. At equilibrium, highly




Figure 5. Minimum population size N_{\min} needed to achieve a local persistence probability $P_{\text{loc}} = 5\%$ and $P_{\text{loc}} = 95\%$ as function of the duration of immunity L for the case of an isolated population and for populations of the metapopulation system. The shaded area denotes the possible population values in our system.

connected and populated patches are steadily infected, while the peripheral ones experience repeated outbreaks on the rare occasions in which infectious travellers arrive from the hubs and find enough susceptible individuals to induce a local outbreak. To characterize this hierarchy, we analyse how local persistence changes across classes of patches and quantify the number of outbreaks each class experiences throughout the duration of the simulated epidemic. Classes of patches are defined by their connectivity, i.e. we focus on nodes having the same number k of neighbouring sub-populations. Owing to the correlation between degree of a patch and its population size, these connectivity classes correspond effectively to population classes. Increasing degree leads therefore to higher local persistence, because of the relation to the population size (figure 6*a* for the low mobility regime). P_{loc} reaches 1 in high-degree nodes for small enough values of the immunity period (i.e. up to $L = 100$ days among the cases shown in the figure 6*a*). In addition, we find that in these conditions hubs consistently experience only one outbreak (figure 6*b*), corresponding to an epidemic that persists in the population after the first seeding event. Low-degree nodes instead experience repeated outbreaks seeded from neighbouring nodes interspersed with local extinctions. For higher values of L (e.g. $L = 150$), the maximum local persistence achieved even in highly connected nodes is smaller than 1 and global persistence is approximately $P_{\text{gl}} = 0.03$. Our results show that when disease persistence occurs, it is driven by repeated outbreaks taking place in different locations. Patches with intermediate degrees have the highest turnover of consecutive epidemics (about three outbreaks measured in the explored timeframe; figure 6*b*). Hubs present fewer outbreaks, as these last longer because of their larger population size and due to the continuous reseeding from neighbouring patches that lead to multiple consecutive waves. Peripheral nodes instead have less outbreaks because, being less reachable, they are not frequently reseeded after local extinctions by other nodes.

3.2.2. Intermediate mobility coupling

For intermediate values of the travelling probability p , the spatial dynamics displays multiple waves followed by an endemic behaviour, if the probability of global persistence is larger than 5% (figure 4*b,c,h*). For smaller persistence (e.g. $P_{\text{gl}} = 0.01$, figure 4*a*), the epidemic goes extinct after an initial wave. Interestingly, the amplitude of the initial wave is determined only by the mobility, regardless of the final persistence or extinction outcome. We explain this by noticing that the first wave refers to the process of spatial invasion from the initially seeded patch, that can be quantified by the global invasion threshold [20]:

$$R_* = (R_0 - 1) \frac{\langle k^2 \rangle - \langle k \rangle^2}{\langle k \rangle^2} \frac{2(R_0 - 1)^2}{\mu R_0^2} p \langle N \rangle, \quad (3.2)$$




Figure 6. (a,c,e) Probability of local persistence in patches with degree k for different values of the immunity period L at $p = 3 \times 10^{-5}$ (top row), 6×10^{-4} (central row) and 10^{-2} (bottom row). The probability is computed as the fraction of outbreaks occurring in a node of degree k lasting at least 2000 time-steps. (b,d,f) Average number of distinct outbreaks experienced during a simulation by a node of degree k for different values of the immunity period L at $p = 3 \times 10^{-5}$ (top row), 6×10^{-4} (central row) and 10^{-2} (bottom row). Statistics is limited to runs where global persistence occurs. All measurements have been performed after a transient time of 2000 steps to focus on the endemic dynamics.

measuring the number of neighbouring patch that a seeded patch can infect. This is the analogue of the basic reproductive number at the sub-population level. Though this estimate was obtained for a SIR dynamics, we consider it here as we focus on the initial spatial invasion following the seeding, before the mechanism of immunity waning becomes important. The invasion condition $R_* > 1$ indeed




Figure 7. (a) Temporal evolution of the fraction of infected individuals (curve) and imported cases (points) in the largest population of the metapopulation system ($k = 91$, $N = 2.5 \times 10^5$) for $p = 6 \times 10^{-4}$ and duration of immunity $L = 320$. (b) Temporal evolution of the fraction of infected individuals (curve) and imported cases (points) in the same patch for $p = 10^{-2}$ and duration of immunity $L = 320$. Both panels show the results of a single stochastic run.

already proved to be a good approximation to define the upper bound of the low mobility regime in equation (3.1). Exploring different points in the intermediate mobility regime characterized by the same value of p ($p = 6 \times 10^{-4}$ for figure 4*a,b,c,h*), we find indeed the same initial invasion, regardless of the duration of the immunity period.

Following the initial wave, dampened oscillations in the number of infected patches are observed at the spatial level, for large enough persistence probability. At the local level, oscillations in the incidence are caused by the waning of immunity and have a frequency that decreases with L and the patch population size, as predicted by the SIRS theory [4]. These local oscillations then interact at the spatial level because of the mobility coupling, translating in multiple waves in the spatial propagation. Those may be the result of travelling waves departing from well-connected nodes [23,24,48,49]. Figure 6*c* indeed shows that if the pathogen is able to persist in the population, after the initial transient, the epidemic has more chances to persist in highly connected nodes. The more peripheral patches experience a sequence of sporadic outbreaks, similarly to the low mobility case (figure 6*d*). Differently from before, here the local persistence is enhanced by the spatial coupling provided by the intermediate mobility. The continuous reseeding by infectious individuals travelling between sub-populations, in a situation of low level of synchronization between local epidemics, fuel the circulation of the infection, hence allowing for a non-zero persistence even at high values of L , corresponding to the maximum discussed in §3.1.

This *rescue effect* was previously introduced in the context of childhood diseases [26–28]. Here, we find that it is driven by the increase in local persistence in highly connected nodes as illustrated in figures 5 and 6. In figure 5, the comparison of this mobility regime with the low mobility one shows that the rescue effect lowers the population threshold needed to have persistence $P_{gl} \geq 0.05$ and $P_{gl} \geq 0.95$. Moreover, comparing the intermediate and low mobility regimes (blue curves in figure 6*a,c*) for the case $L = 150$, we see that while local persistence is not achieved when patches are fairly isolated, this is not the case for a large class of nodes as soon as mobility increases.

3.2.3. High mobility coupling

To characterize the unfolding of the epidemic in the high mobility regime, we consider two points of figure 4*e* having the same travelling probability $p = 10^{-2}$, and different lengths of the immunity period, namely $L = 150$ and $L = 250$, corresponding to $P_{gl} = 1.0$ and $P_{gl} = 0.60$, respectively. The epidemic dynamics in these cases shows an oscillatory behaviour followed by an endemic equilibrium (figure 4*f,i*), similar to what was previously discussed in the intermediate mobility regime. Here, frequencies of oscillations are, however, higher. This may be induced by an increase in synchronization among patches, enabled by the stronger level of spatial coupling [31]. Larger travelling fluxes lead, indeed, to almost all patches being infected during the first wave (e.g. $D/V \simeq 1$ for $p = 10^{-2}$, figure 4*i*, compared with $D/V \simeq 0.6$ for $p = 6 \times 10^{-4}$, figure 4*h*, for the same value of L). Over time, these fluxes guarantee a large degree of seeding across patches along with increased mixing. For $L \lesssim 360$ and in a situation of global persistence, local persistence is found to be much higher across degree classes compared with intermediate mobility—see e.g. curves in figure 6*e* versus figure 6*c* for the same value of L and consistently the values of N_{\min} in figure 5 corresponding to the $p = 10^{-2}$ case. Spatial heterogeneities are therefore progressively reduced, with a larger number of patches behaving similarly and epidemic differences across patches almost disappearing.

If persistence is enhanced locally for all patches, the resulting more synchronized system is, however, more prone to extinction of the epidemic for large enough immunity periods. This is the same type of phenomenon observed for measles in [26–28], i.e. a decrease in spatial heterogeneity reduces the effectiveness of the rescue effect by enhancing the level of synchronization. To test this hypothesis, we compared the local and importation dynamics occurring in the most connected patch in the intermediate and high mobility regimes. We considered a single stochastic run to highlight the critical role of stochastic effects within this mechanism (figure 7). Immunity is maintained equal in both scenarios, for comparison. When $p = 6 \times 10^{-4}$ (intermediate mobility, figure 7*a*), importation of cases from neighbouring patches, desynchronized with respect to the local dynamics, allows the epidemic to survive the first wave and then quickly reach an endemic equilibrium, i.e. the rescue effect referred to before. If mobility is higher ($p = 0.01$, figure 7*b*), importations are synchronized with the local dynamics, i.e. a peak of importations is occurring during the peak of the local epidemic wave, and both profiles fade out at the same time. In this way, there is no importation from outside the patch that could help sustain or relaunch the outbreak in the patch, and the epidemic goes extinct.

4. Conclusion

As has been previously discussed in the literature, the epidemiological mechanism of waning of immunity alters the fate of an emerging infection and its chance to persist in the population. Here, we have characterized in depth this effect once a spatially structured population with varying coupling is considered. By using a stochastic individual-based metapopulation model that explicitly accounts for travelling fluxes among sub-populations, we have found that there are three distinct mobility regimes. Our results show that, when mobility is low, the persistence conditions for the disease in the metapopulation system are the same as if the patches were isolated. When mobility is very high the epidemic behaves as if the whole metapopulation were well mixed. Eventually, at intermediate levels of mobility, the coupling between sub-populations creates rescue effects due to non-synchronized epidemics ongoing in the patches that maximize disease persistence. These effects are able to reseed the epidemics locally once it became extinct.

Different persistence outcomes are associated with a variety of short-term dynamical behaviours. These are due to the interplay between local replenishment of susceptible individuals and the level of epidemic synchronization across patches. For a very rapid waning of immunity, persistence occurs regardless of the level of mobility coupling, with local endemic equilibria allowing for the epidemic to propagate from one patch to another even at very low travelling probabilities. For longer

durations of the immunity period, the probability of having local persistence is shaped by the high heterogeneity in the population distribution (large patches may be above the critical size needed for local persistence, while small ones may not) and the rescue effect. These two mechanisms are important at intermediate levels of travelling coupling and are responsible for the peak of persistence observed in this regime.

Our findings are similar in many aspects to the ones obtained for the case of measles and, more in general, for an SIR model with vital dynamics and external introductions [23,24,27,28,30,31,48,50–52]. This exemplifies that different mechanisms of susceptible replenishment can lead to common dynamical features. Distinct and apparently contradicting behaviours have been shown in previous studies [28–31], highlighting the important role that mobility can have in triggering these mechanisms. Here, we have provided a detailed description of the epidemic dynamics for the case of a rapidly spreading acute infection with waning of immunity, where the main statistical features of human space distribution and mobility are accounted for in a realistic way. We showed that the main result, i.e. that persistence is maximized at intermediate mobility couplings, is valid for different levels of spatial fragmentation and also for a homogeneous distribution of population and connectivity.

In order to better focus the analysis on the interplay between mobility and time scale of waning of immunity, we considered throughout the manuscript a single value of R_0 . We expect different values of R_0 yielding the same qualitative picture with an increase in the parameter determining a shift of the persistence peak toward smaller values of p as predicted by equation (3.1). We also disregarded ingredients known to impact infectious disease dynamics in many real cases. In particular, we assumed no seasonality in transmission and no multi-strain interference. These two ingredients are known to alter the spread of diseases like influenza and respiratory syncytial virus, thus they are usually included in realistic data-driven modelling of these infections [2,7,53]. The inclusion of seasonality in the studies of childhood diseases and its interplay with demography showed the emergence of annual and biannual cycles in the dynamics of the disease [23,24,51]. In the systematic exploration conducted here for the SIRS case, accounting for these ingredients would be complicated by the large increase in the complexity of the dynamics, the number of parameters and the variety of possible outcomes in terms of persistence, transient dynamics and equilibrium state [21,22,29,54–56], thus hindering a clear theoretical understanding. The chosen approximations may, however, limit the applicability of our work directly to real-case situations, such as the case of the emerging influenza strains [2,57–59].

Other aspects could alter the results, such as human response and intervention to the epidemic—e.g. vaccination, case isolation and travel restrictions. In particular, phenomena like travel bans or spontaneous reductions of travel often occur during the initial stage of an outbreak as a consequence of the public concern about the emerging infection. They were shown, however, to have limited impact on the early stage invasion [19], given that reductions of orders of magnitude needed to significantly affect spatial spread are unfeasible in reality [60,61]. For the same reason we argue that their effect would be minimal also on pathogen persistence, unless the system is in the intermediate mobility regime, where probability of persistence is more sensitive to the level of spatial coupling (figure 2). Vaccination, on the other hand, by boosting the level of immunity in the population increases the chance of pathogen extinction and may profoundly alter the spreading dynamics, affecting the frequency of epidemic waves [23]. For the case of rapidly mutating pathogens the quantification of vaccine efficacy and its impact on disease eradication is complicated by the fact that immunity caused by vaccination also decays in time when novel virus variants emerge, and vaccine match with the circulating pathogen is often compromised due to the difficulty in anticipating virus mutation in the vaccine design and production [58].

Starting from the complete characterization of the epidemic dynamics proposed here, we plan to address these points in the future in order to better understand the epidemic dynamics resulting from a complex interplay of different factors.

Authors' contributions. S.M., C.P., V.C. and Y.M. conceived the study with A.N.S.H. contributing at the initial phase. A.N.S.H. carried out preliminary numerical analysis and A.A. performed the final numerical simulations. A.N.S.H. participated in the analysis of preliminary results and A.A., S.M., C.P., V.C. and Y.M. analysed the final results. All authors contributed to writing the manuscript and approved the final version.

Competing interests. The authors declare no competing interests.

Funding. S.M. acknowledges support from the Juan de la Cierva Program, Spain. This work was partially supported by the EC-Health Contract PREDEMICS (No. 278433) and the ANR Contract HARMSFLU (No. ANR-12-MONU-0018) to V.C.; the Government of Aragón, Spain through a grant to the group FENOL to Y.M.; the MINECO and FEDER

funds (grant FIS2014-55867-P) to Y.M.; the European Commission FET-Proactive Project Multiplex (grant no. 317532) to Y.M.; the Ciência sem Fronteiras programme—PDE/CNPq (No. 245936/2012-2) to A.N.S.H.

Acknowledgements. A.N.S.H. gratefully acknowledges L. H. G. Tizei for insightful discussions.

References

- Keeling MJ, Rohani P. 2008 *Modeling infectious diseases in humans and animals*. Princeton, NJ: Princeton University Press.
- Zhang XS, De Angelis D, White PJ, Charlett A, Pebody RG, McCauley J. 2013 Co-circulation of influenza A virus strains and emergence of pandemic via reassortment: the role of cross-immunity. *Epidemics* **5**, 20–33. (doi:10.1016/j.epidem.2012.10.003)
- Weber A, Weber M, Milligan P. 2001 Modeling epidemics caused by respiratory syncytial virus (RSV). *Math. Biosci.* **172**, 95–113. (doi:10.1016/S0025-5564(01)00066-9)
- Pease CM. 1987 An evolutionary epidemiological mechanism, with applications to type A influenza. *Theor. Popul. Biol.* **31**, 422–452. (doi:10.1016/0040-5809(87)90014-1)
- Gomes MGM, White LJ, Medley GF. 2004 Infection, reinfection, and vaccination under suboptimal immune protection: epidemiological perspectives. *J. Theor. Biol.* **228**, 539–549. (doi:10.1016/j.jtbi.2004.02.015)
- Bansal S, Meyers LA. 2012 The impact of past epidemics on future disease dynamics. *J. Theor. Biol.* **309**, 176–184. (doi:10.1016/j.jtbi.2012.06.012)
- Truscott J, Fraser C, Cauchemez S, Meeyai A, Hinsley W, Donnelly CA, Ghani A, Ferguson N. 2011 Essential epidemiological mechanisms underpinning the transmission dynamics of seasonal influenza. *J. R. Soc. Interface* **9**, 304–312. (doi:10.1098/rsif.2011.0309)
- Näsell I. 2013 The influence of immunity loss on persistence and recurrence of endemic infections. *Bull. Math. Biol.* **75**, 2079–2092. (doi:10.1007/s11538-013-9882-x)
- Clancy D, Mendy ST. 2009 The effect of waning immunity on long-term behaviour of stochastic models for the spread of infection. *J. Math. Biol.* **61**, 527–544. (doi:10.1007/s00285-009-0313-4)
- Yuan HY, Koelle K. 2013 The evolutionary dynamics of receptor binding avidity in influenza A: a mathematical model for a new antigenic drift hypothesis. *Phil. Trans. R. Soc. B* **368**, 20120204. (doi:10.1098/rstb.2012.0204)
- Andreasen V, Lin J, Levin SA. 1997 The dynamics of cocirculating influenza strains conferring partial cross-immunity. *J. Math. Biol.* **35**, 825–842. (doi:10.1007/s002850050079)
- Shaman J, Pitzer VE, Viboud C, Grenfell BT, Lipsitch M. 2010 Absolute humidity and the seasonal onset of influenza in the continental United States. *PLoS Biol.* **8**, e1000316. (doi:10.1371/journal.pbio.1000316)
- Dushoff J, Plotkin JB, Levin SA, Earn DJD. 2004 Dynamical resonance can account for seasonality of influenza epidemics. *Proc. Natl Acad. Sci. USA* **101**, 16 915–16 916. (doi:10.1073/pnas.0407293101)
- Li D, Cui J, Liu M, Liu S. 2015 The evolutionary dynamics of stochastic epidemic model with nonlinear incidence rate. *Bull. Math. Biol.* **77**, 1705–1743. (doi:10.1007/s11538-015-0101-9)
- Casagrandi R, Bolzoni L, Levin SA, Andreasen V. 2006 The SIRC model and influenza A. *Math. Biosci.* **200**, 152–169. (doi:10.1016/j.mbs.2005.12.029)
- Asaduzzaman SM, Ma J, den Driessche P. 2015 The coexistence or replacement of two subtypes of influenza. *Math. Biosci.* **270**, 1–9. (doi:10.1016/j.mbs.2015.09.006)
- Balcan D, Vespignani A. 2011 Phase transitions in contagion processes mediated by recurrent mobility patterns. *Nat. Phys.* **7**, 581–586. (doi:10.1038/nphys1944)
- Belik V, Geisel T, Brockmann D. 2011 Natural human mobility patterns and spatial spread of infectious diseases. *Phys. Rev. X* **1**, 011001. (doi:10.1103/PhysRevX.1.011001)
- Meloni S, Perra N, Arenas A, Gómez S, Moreno Y, Vespignani A. 2011 Modeling human mobility responses to the large-scale spreading of infectious diseases. *Sci. Rep.* **1**, 62. (doi:10.1038/srep00062)
- Colizza V, Vespignani A. 2008 Epidemic modeling in metapopulation systems with heterogeneous coupling pattern: theory and simulations. *J. Theor. Biol.* **251**, 450–467. (doi:10.1016/j.jtbi.2007.11.028)
- Poletto C, Meloni S, Colizza V, Moreno Y, Vespignani A. 2013 Host mobility drives pathogen competition in spatially structured populations. *PLoS Comput. Biol.* **9**, e1003169. (doi:10.1371/journal.pcbi.1003169)
- Poletto C, Meloni S, Van Metre A, Colizza V, Moreno Y, Vespignani A. 2015 Characterising two-pathogen competition in spatially structured environments. *Sci. Rep.* **5**, 7895. (doi:10.1038/srep07895)
- Grenfell BT, Bjørnstad ON, Kappey J. 2001 Travelling waves and spatial hierarchies in measles epidemics. *Nature* **414**, 716–723. (doi:10.1038/414716a)
- Marguta R, Parisi A. 2015 Impact of human mobility on the periodicities and mechanisms underlying measles dynamics. *J. R. Soc. Interface* **12**, 20141317. (doi:10.1098/rsif.2014.1317)
- Bartlett MS. 1957 Measles periodicity and community size. *J. R. Stat. Soc. Ser. A* **120**, 48–70. (doi:10.2307/2342553)
- Bolker B, Grenfell B. 1995 Space, persistence and dynamics of measles epidemics. *Phil. Trans. R. Soc. Lond. B* **348**, 309–320. (doi:10.1098/rstb.1995.0070)
- Bolker BM, Grenfell BT. 1996 Impact of vaccination on the spatial correlation and persistence of measles dynamics. *Proc. Natl Acad. Sci. USA* **93**, 12 648–12 653. (doi:10.1073/pnas.93.22.12648)
- Keeling MJ. 2000 Metapopulation moments: coupling, stochasticity and persistence. *J. Anim. Ecol.* **69**, 725–736. (doi:10.1046/j.1365-2656.2000.00430.x)
- Rozhnova G, Nunes A, McKane AJ. 2014 Impact of commuting on disease persistence in heterogeneous metapopulations. *Ecol. Complex.* **19**, 124–129. (doi:10.1016/j.ecocom.2014.06.001)
- Jesse M, Heesterbeek H. 2011 Divide and conquer? Persistence of infectious agents in spatial metapopulations of hosts. *J. Theor. Biol.* **275**, 12–20. (doi:10.1016/j.jtbi.2011.01.032)
- Hagenaars TJ, Donnelly CA, Ferguson NM. 2004 Spatial heterogeneity and the persistence of infectious diseases. *J. Theor. Biol.* **229**, 349–359. (doi:10.1016/j.jtbi.2004.04.002)
- Levins R. 1969 Some demographic and genetic consequences of environmental heterogeneity for biological control. *Bull. Entomol. Soc. Am.* **15**, 237–240. (doi:10.1093/besa/15.3.237)
- Hanski I, Gaggiotti OE. 2004 *Ecology, genetics, and evolution of metapopulations*. New York, NY: Academic Press.
- Lloyd AL, May RM. 1996 Spatial heterogeneity in epidemic models. *J. Theor. Biol.* **179**, 1–11. (doi:10.1006/jtbi.1996.0042)
- Balcan D, Colizza V, Gonçalves B, Hu H, Ramasco JJ, Vespignani A. 2009 Multiscale mobility networks and the spatial spreading of infectious diseases. *Proc. Natl Acad. Sci. USA* **106**, 21 484–21 489. (doi:10.1073/pnas.0906910106)
- Simini F, González MC, Maritan A, Barabási AL. 2012 A universal model for mobility and migration patterns. *Nature* **484**, 96–100. (doi:10.1038/nature10856)
- Chowell G, Hyman JM, Eubank S, Castillo-Chavez C. 2003 Scaling laws for the movement of people between locations in a large city. *Phys. Rev. E* **68**, 066102. (doi:10.1103/PhysRevE.68.066102)
- Barrat A, Barthélemy M, Pastor-Satorras R, Vespignani A. 2004 The architecture of complex weighted networks. *Proc. Natl Acad. Sci. USA* **101**, 3747–3752. (doi:10.1073/pnas.0400087101)
- Catanzaro M, Boguná M, Pastor-Satorras R. 2005 Generation of uncorrelated random scale-free networks. *Phys. Rev. E* **71**, 027103. (doi:10.1103/PhysRevE.71.027103)
- Erdős P, Rényi Á. 1959 On random graphs I. *Publicationes Mathematicae (Debrecen)*. **6**, 290–297.
- Cross PC, Lloyd-Smith JO, Johnson PLF, Getz WM. 2005 Duelling timescales of host movement and disease recovery determine invasion of disease in structured populations. *Ecol. Lett.* **8**, 587–595. (doi:10.1111/j.1461-0248.2005.00760.x)
- Ball F, Mollison D, Scalia-Tomba G. 1997 Epidemics with two levels of mixing. *Ann. Appl. Prob.* **7**, 46–89. (doi:10.1214/aoap/1034625252)
- Poletto C, Tizzoni M, Colizza V. 2013 Human mobility and time spent at destination: impact on spatial epidemic spreading. *J. Theor. Biol.* **338**, 41–58. (doi:10.1016/j.jtbi.2013.08.032)
- Colizza V, Vespignani A. 2007 Invasion threshold in heterogeneous metapopulation networks. *Phys. Rev. Lett.* **99**, 148701. (doi:10.1103/PhysRevLett.99.148701)
- Apolloni A, Poletto C, Ramasco JJ, Jensen P, Colizza V. 2014 Metapopulation epidemic models with heterogeneous mixing and travel behaviour. *Theor. Biol. Med. Modell.* **11**, 3. (doi:10.1186/1742-4682-11-3)

46. Wearing HJ, Rohani P. 2009 Estimating the duration of pertussis immunity using epidemiological signatures. *PLoS Pathogens* **5**, e1000647. (doi:10.1371/journal.ppat.1000647)
47. Brooks-Pollock E, Keeling M. 2009 Herd size and bovine tuberculosis persistence in cattle farms in Great Britain. *Prev. Vet. Med.* **92**, 360–365. (doi:10.1016/j.prevetmed.2009.08.022)
48. Grenfell B. 1998 Cities and villages: infection hierarchies in a measles metapopulation. *Ecol. Lett.* **1**, 63–70. (doi:10.1046/j.1461-0248.1998.00016.x)
49. Bjørnstad ON, Ims RA, Lambin X. 1999 Spatial population dynamics: analyzing patterns and processes of population synchrony. *Trends Ecol. Evol.* **14**, 427–432. (doi:10.1016/S0169-5347(99)01677-8)
50. Lande R, Engen S, Sæther BE. 1998 Extinction times in finite metapopulation models with stochastic local dynamics. *Oikos* **83**, 383–389. (doi:10.2307/3546853)
51. Marguta R, Parisi A. 2016 Periodicity, synchronization and persistence in pre-vaccination measles. *J. R. Soc. Interface* **13**, 20160258. (doi:10.1098/rsif.2016.0258)
52. Rohani P, Earn DJD, Grenfell BT. 1999 Opposite patterns of synchrony in sympatric disease metapopulations. *Science* **286**, 968–971. (doi:10.1126/science.286.5441.968)
53. Pitzer VE, Viboud C, Alonso WJ, Wilcox T, Metcalf CJ, Steiner CA, Haynes AK, Grenfell BT. 2015 Environmental drivers of the spatiotemporal dynamics of respiratory syncytial virus in the United States. *PLoS Pathog.* **11**, e1004591. (doi:10.1371/journal.ppat.1004591)
54. Roche B, Drake JM, Rohani P. 2011 An agent-based model to study the epidemiological and evolutionary dynamics of Influenza viruses. *BMC Bioinform.* **12**, 87. (doi:10.1186/1471-2105-12-87)
55. Ballesteros S, Vergu E, Cazes B. 2009 Influenza A gradual and epochal evolution: insights from simple models. *PLoS ONE* **4**, e7426. (doi:10.1371/journal.pone.0007426)
56. Buscarino A, Fortuna L, Frasca M, Rizzo A. 2014 Local and global epidemic outbreaks in populations moving in inhomogeneous environments. *Phys. Rev. E* **90**, 042813. (doi:10.1103/PhysRevE.90.042813)
57. Boëlle PY, Ansart S, Cori A, Valleron AJ. 2011 Transmission parameters of the A/H1N1 (2009) influenza virus pandemic: a review. *Influenza Other Respiratory Viruses* **5**, 306–316. (doi:10.1111/j.1750-2659.2011.00234.x)
58. Bedford T, Suchard MA, Lemey P, Dudas G, Gregory V, Hay AJ, McCauley JW, Russell CA, Smith DJ, Rambaut A. 2014 Integrating influenza antigenic dynamics with molecular evolution. *eLife* **3**, e01914. (doi:10.7554/eLife.01914)
59. He D, Lui R, Wang L, Tse CK, Yang L, Stone L. 2015 Global spatio-temporal patterns of influenza in the post-pandemic era. *Sci. Rep.* **5**, 11013. (doi:10.1038/srep11013)
60. Hollingsworth TD, Ferguson NM, Anderson RM. 2006 Will travel restrictions control the international spread of pandemic influenza? *Nat. Med.* **12**, 497–499. (doi:10.1038/nm0506-497)
61. Poletto C *et al.* 2014 Assessing the impact of travel restrictions on international spread of the 2014 West African Ebola epidemic. *Eurosurveillance* **19**. (doi:10.2807/1560-7917.ES2014.19.42.20936)