

manuscript No.
(will be inserted by the editor)

G. Brida · M. Bondani
I. P. Degiovanni · M. Genovese
M. G. A. Paris · I. Ruo Berchera
V. Schettini

On the discrimination between classical and quantum states

the date of receipt and acceptance should be inserted later

Abstract With the purpose to introduce a useful tool for researches concerning foundations of quantum mechanics and applications to quantum technologies, here we study three quantumness quantifiers for bipartite optical systems: one based on sub-shot-noise correlations, one related to antibunch-

This work has been supported by SanPaolo Foundation, Quantum Candela EU Project

G. Brida
Istituto Nazionale di Ricerca Metrologica, I-10135, Torino, Italia
E-mail: g.brida@inrim.it

M. Bondani
National Laboratory for Ultrafast and Ultraintense Optical Science,
C.N.R.-I.N.F.M., I-22100 Como, Italia
C.N.I.S.M., U.d.R. Como, I-22100 Como, Italia
E-mail: maria.bondani@uninsubria.it

I. P. Degiovanni
Istituto Nazionale di Ricerca Metrologica, I-10135, Torino, Italia
E-mail: i.degiovanni@inrim.it

M. Genovese
Istituto Nazionale di Ricerca Metrologica, I-10135, Torino, Italia
E-mail: m.genovese@inrim.it

M. G. A. Paris
Dipartimento di Fisica, Università degli Studi di Milano, I-20133 Milano, Italia
C.N.I.S.M., U.d.R. Milano Università, I-20133 Milano, Italia
I.S.I. Foundation, I-10133 Torino, Italia
E-mail: matteo.paris@fisica.unimi.it

I. Ruo Berchera
Istituto Nazionale di Ricerca Metrologica, I-10135, Torino, Italia
E-mail: i.ruoberchera@inrim.it

V. Schettini
Istituto Nazionale di Ricerca Metrologica, I-10135, Torino, Italia
E-mail: v.schettini@inrim.it

ing and one springing from entanglement determination. The specific cases of parametric down conversion seeded by thermal, coherent and squeezed states are discussed in detail.

Keywords entanglement, macro-objectification

PACS 42.50.Dv · 03.67.Mn

1 Introduction

The discrimination between quantum and classical states [1, 2, 3, 4, 5], beyond its very important and deep conceptual relevance, has also recently received much attention due to the development of quantum technologies. On the one hand, it represents a fundamental point for the studies concerning the transition between quantum and classical world, one of the most intriguing research sectors in the foundations of physics; on the other hand it is a tool of the utmost importance when comparing results that can be achieved with quantum and classical protocols.

These studies have concerned various physical systems [1] ranging from quantum optical states [4] to mesons [6] or solid state devices [7] and, recently, have pointed to the need of an operational approach linked to measurements schemes [8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25]. Considering the experimental interest in the frame of quantum optics [25, 26, 27, 28], in a recent paper [29] we have considered three different “quantumness” quantifiers applied to the study of quantum-classical transition in a thermal seeded parametric down conversion (PDC): a work that, a part its specific application, has a more general interest since it presents an idea that can find generalization to various physical systems.

Here we want to extend this first study by considering, as a further example, the application of these three quantifiers to PDC seeded by coherent and squeezed vacuum states, comparing them with what we achieved for thermal seeded case. This allows a more general understanding of the hierarchy of these three quantifiers that emerged in [29] and, due to the easy realizability of these states, paves the way toward a general experimental test of these theoretical results.

The structure of the paper is the following. After an introduction to seeded PDC (section 2), we consider the sub-shot noise measurement (section 3), Lee’s criterion (section 4) and entanglement (section 5). Finally, a general discussion of the results is presented in section 6.

2 Introduction to seeded parametric down-conversion

The evolution of a quantum system induced by the interaction Hamiltonian describing the PDC process for a single pair of coupled modes is described by the unitary operator $U = \exp(i\kappa a_A a_B + h.c.)$, where $\kappa = |\kappa|e^{i\varphi}$ is the coupling constant and a_A and a_B are the annihilation operators for photons on modes A and B, respectively. We consider the PDC process seeded by two single mode input states $\rho_{in} = \rho_A \otimes \rho_B$. In particular as seeds fields

on both A- and B- modes we consider the three simplest Gaussian states, namely thermal states, coherent states, and vacuum squeezed states. For the thermal case (T) the input state of the single channel is a mixed incoherent superposition.

$$\rho_j^{(T)} = \sum_{n=0}^{\infty} P_j(n) |n\rangle_j \langle n|, \quad (1)$$

where $j = A, B$ and $|n\rangle_j$ denotes the Fock number basis for the single mode of the j -arm, the thermal probability distribution of the input being $P_j(n) = \mu_j^n (1 + \mu_j)^{-n-1}$, where μ_j is the average photon number.

In the case of coherent seeding (C) the state in the single channel is obtained by the action of the displacement operator $D(\alpha_j) = e^{i(\alpha_j a_j + \alpha_j^* a_j^\dagger)}$ on the vacuum state

$$\rho_j^{(C)} = D(\alpha_j) |0\rangle_j \langle 0| D^\dagger(\alpha_j) \quad (2)$$

with $\alpha_j = \sqrt{M_j} e^{i\gamma_j}$ is the complex displacement parameters and M_j represents the mean number of photon of the state.

For the squeezed vacuum seeds (S) the input state of the single channel is given by

$$\rho_j^{(S)} = S(\xi_j) |0\rangle_j \langle 0| S^\dagger(\xi_j) \quad (3)$$

where the squeezing operator is $S(\xi_j) = e^{i(\xi_j a_j^2 + \xi_j^* a_j^{\dagger 2})}$, $\xi_j = |\xi_j| e^{i\zeta_j}$ being complex parameters.

The density matrix at the output of the PDC interaction is given by

$$\rho_{out} = U \rho_{in} U^\dagger. \quad (4)$$

Conversely in the interaction picture the output field modes are given by $A_j = U^\dagger a_j U$, *i.e*

$$A_j = \sqrt{N+1} a_j + e^{i\varphi} \sqrt{N} a_{j'}^\dagger \quad (j, j' = A, B, j \neq j') \quad (5)$$

where $N = \sinh^2 |\kappa|$ is the mean number of photons of the PDC spontaneous emission.

The first moments of the photon number distribution in the case of thermal seeds are [31]

$$\begin{aligned} \langle n_A^{(T)} \rangle &= \mu_A + N(1 + \mu_A + \mu_B) \\ \langle n_B^{(T)} \rangle &= \mu_B + N(1 + \mu_A + \mu_B) \end{aligned} \quad (6)$$

where, $n_j = a_j^\dagger a_j$, $\langle O \rangle = \text{Tr}[O \rho_{out}]$.

When the process is seeded by coherent fields we have:

$$\begin{aligned} \langle n_A^{(C)} \rangle &= M_A + N(1 + M_A + M_B) + 2\sqrt{N(N+1)}\sqrt{M_A M_B} \cos(\gamma_A + \gamma_B - \varphi) \\ \langle n_B^{(C)} \rangle &= M_B + N(1 + M_A + M_B) + 2\sqrt{N(N+1)}\sqrt{M_A M_B} \cos(\gamma_A + \gamma_B - \varphi) \end{aligned} \quad (7)$$

Contrary to the thermal case, here the intensity of the fields is partially modulated by the phase value of the seeding fields with respect to the phase induced by the PDC process when both the seeds are nonzero (i.e. $M_A, M_B \neq 0$). Eventually when vacuum squeezed input beams are considered, we obtain

$$\begin{aligned}\langle n_A^{(S)} \rangle &= N_A + N(1 + N_A + N_B) \\ \langle n_B^{(S)} \rangle &= N_B + N(1 + N_A + N_B),\end{aligned}\tag{8}$$

where $N_j = \sinh^2 |\xi_j|$ is the average number of photons of the input state in the single channel j , whereas, surprisingly, the phases of the squeezing operators do not play any role in the photon number. Notice that the case of vacuum inputs, $\rho_{in} = |0\rangle\langle 0|_A \otimes |0\rangle\langle 0|_B$, corresponds to spontaneous PDC, *i.e.* to the generation of twin-beam.

3 Sub-Shot Noise Measurement

The shot-noise limit (SNL) in any photodetection process is defined as the lowest level of noise that can be obtained by using semiclassical states of light that is, Glauber coherent states. If one measures the photon numbers in two beams and evaluates their difference, the SNL is the lower limit of noise that can be reached when the beams are classically correlated. On the other hand, when intensity correlations below the SNL are observed, we have a genuine nonclassical effect. We consider a simple measurement scheme where A and B single mode beams at the output of the PDC interaction are individually measured by direct detection (considering in this section ideal detectors with unitary efficiency [30]). The resulting A and B photon counts, which are correlated, are subtracted from each other to demonstrate quantum noise reduction in the difference of photon counts. In order to observe a violation of the SNL we must have

$$\langle [\Delta(n_A - n_B)]^2 \rangle < \langle n_A \rangle + \langle n_B \rangle\tag{9}$$

where $\langle [\Delta(n_A - n_B)]^2 \rangle$ is the variance of this difference, and $\langle n_A \rangle + \langle n_B \rangle$ is the SNL, *i.e.* the quantity that would be obtained for uncorrelated coherent beams.

In particular, the condition in Eq. (9) reduces to

$$N > \frac{(\mu_A^2 + \mu_B^2)}{2(1 + \mu_A + \mu_B)}\tag{10}$$

for the thermal seeds.

In the case of coherent seeds if the phases satisfy $\cos(\gamma_A + \gamma_B - \varphi) \geq 0$ the condition in Eq. (9) is always fulfilled irrespective of the value of N , M_A , M_B , while if $\cos(\gamma_A + \gamma_B - \varphi) < 0$ the condition is fulfilled only when

$$N > \frac{4M_A M_B \cos^2(\gamma_A + \gamma_B - \varphi)}{1 + 2(M_A + M_B) + (M_A + M_B)^2 - 4M_A M_B \cos^2(\gamma_A + \gamma_B - \varphi)}.\tag{11}$$

In the case of squeezed vacuum seeds the condition is

$$N > \frac{N_A(1 + 2N_A) + N_B(1 + 2N_B)}{2(1 + N_A + N_B)}. \quad (12)$$

It is interesting to notice that, in the case of thermal and squeezed input state, there always exists a threshold between the sub-shot-noise and the classical regime, which can be explored by controlling the intensities of the seeds. The behavior of the coherent case is different because, upon properly adjusting the phases, the sub-shot noise condition holds whatever the intensities of the seeds. In an experiment in which the phases of seeds γ_A, γ_B and that of PDC process φ are not locked, one expects, on average, a null value of the cosine in Eq. (11) and therefore a permanent SSN condition. It is helpful to define a parameter, \mathcal{P}_{SSN} quantifying the amount of violation of the SNL

$$\mathcal{P}_{SSN} = 1 - \frac{\langle [\Delta(n_A - n_B)]^2 \rangle}{\langle n_A \rangle + \langle n_B \rangle}. \quad (13)$$

$\mathcal{P}_{SSN} = 0$ corresponds to the SNL, and the sub-shot noise condition corresponds to $0 < \mathcal{P}_{SSN} \leq 1$. For the state ρ_{out} , in the case of thermal seeds we have

$$\mathcal{P}_{SSN}^{(T)} = \frac{2 \mu_{PDC}(1 + \mu_A + \mu_B) - \mu_A^2 - \mu_B^2}{2 \mu_{PDC}(1 + \mu_A + \mu_B) + \mu_A + \mu_B}, \quad (14)$$

thus the maximal violation of SNL is achieved by the twin-beam ($\mu_A = \mu_B = 0$), and by increasing the magnitude of, at least, one of the seeding field the SNL is reached.

For coherent input beams the amount of violation is

$$\mathcal{P}_{SSN}^{(C)} = \frac{2N(1 + M_A + M_B) + 4\sqrt{N(N+1)}\sqrt{M_A M_B} \cos(\gamma_A + \gamma_B - \varphi)}{2N(1 + M_A + M_B) + 4\sqrt{N(N+1)}\sqrt{M_A M_B} \cos(\gamma_A + \gamma_B - \varphi) + M_A + M_B}, \quad (15)$$

and also in this case the limit value of 1 is reached again by the twin-beams in the spontaneous emission ($M_A = M_B = 0$). The SNL threshold $\mathcal{P}_{SSN}^{(C)} = 0$ is obtained when the numerator of Eq. (15) is zero, leading to the solutions presented in Eq.(11).

Finally, for the squeezed beams the parameter is

$$\mathcal{P}_{SSN}^{(S)} = \frac{2N(1 + N_A + N_B) - 2N_A(1 + N_A) - 2N_B(1 + N_B)}{2N(1 + N_A + N_B) + N_A + N_B}. \quad (16)$$

We notice that for all the cases, it can be shown that $\langle [\Delta(n_A - n_B)]^2 \rangle$ is equal to the sum of the mean fluctuation of the two input seeding states. Therefore, \mathcal{P}_{SSN} always assumes the form

$$\mathcal{P}_{SSN} \equiv 1 - \frac{\langle [\Delta(n_A - n_B)]^2 \rangle}{\langle n_A \rangle + \langle n_B \rangle} = \frac{\langle n_A \rangle + \langle n_B \rangle - \langle [\Delta n_A]^2 \rangle_{\rho_{in}} - \langle [\Delta n_B]^2 \rangle_{\rho_{in}}}{\langle n_A \rangle + \langle n_B \rangle}, \quad (17)$$

where, with obvious notation, $\langle O \rangle_{\rho_{in}} = \text{Tr}[O\rho_{in}]$.

4 Lee's Criterion

Another interesting criterion of nonclassicality was derived by Lee [32,33], and it is the two-mode generalization of the well known nonclassicality criterion for single mode beam $\langle n(n-1) \rangle - \langle n \rangle^2 < 0$ [34]. The Lee's criterion states that a bipartite system is nonclassical if the inequality

$$\langle n_A(n_A - 1) \rangle + \langle n_B(n_B - 1) \rangle - 2\langle n_A n_B \rangle < 0 \quad (18)$$

is satisfied. It is noteworthy to observe that the "Lee's nonclassicality" corresponding to Eq. (18), implies the negativity of the Glauber-Sudarshan P-function [32,33]. Once we consider the state ρ_{out} , the condition in Eq. (18) for seeded PDC is achieved when

$$N > \frac{\mu_A^2 + \mu_B^2 - \mu_A \mu_B}{(1 + \mu_A + \mu_B)}, \quad (19)$$

while in the case of coherent seeds, we have that, when the phases satisfy $\cos(\gamma_A + \gamma_B - \varphi) \geq 0$, $N > N_-$, while when $\cos(\gamma_A + \gamma_B - \varphi) < 0$, $N > N_+$ with

$$N_{\pm} = \frac{4g \cos^2 r + ab \pm 2[g \cos^2 r (4g \cos^2 r + 2ab + a^2)]^{1/2}}{2[b^2 - 4g \cos^2 r]} \quad (20)$$

and $g = M_A M_B$, $a = (M_A - M_B)^2$, $b = 1 + M_A + M_B$, $r = \gamma_A + \gamma_B - \varphi$.

For the squeezed thermal seeds we obtain

$$N > \frac{N_A + N_B + 3N_A^2 + 3N_B^2 - 2N_A N_B}{2(1 + N_A + N_B)}. \quad (21)$$

It is noteworthy to observe that the Lee condition is stricter than the sub-SNL: always exists a threshold between classicality and nonclassicality for the Lee's criterion.

Analogously to the sub-SNL case we define a parameter quantifying the amount of violation of the classicality bound as

$$\mathcal{P}_{Lee} = 1 - \frac{\langle [\Delta(n_A - n_B)]^2 \rangle + (\langle n_A \rangle - \langle n_B \rangle)^2}{\langle n_A \rangle + \langle n_B \rangle}. \quad (22)$$

$\mathcal{P}_{SSN} = 0$ corresponds to the bound of the Lee nonclassicality region, and the Lee nonclassicality condition corresponds to $0 < \mathcal{P}_{Lee} \leq 1$. For the thermal seeds we obtain

$$\mathcal{P}_{Lee}^{(T)} = 2 \frac{N(1 + \mu_A + \mu_B) - \mu_A^2 - \mu_B^2 + \mu_A \mu_B}{2N(1 + \mu_A + \mu_B) + \mu_A + \mu_B}. \quad (23)$$

Thus, the maximal violation of Lee's criterion ($\mathcal{P}_{Lee} = 1$) is achieved by the twin-beam ($\mu_A = \mu_B = 0$), and by increasing the magnitude of, at least, one of the seeding field the classicality bound is reached.

For coherent input beams, the Lee parameter is

$$\mathcal{P}_{Lee}^{(C)} = \frac{2N(1 + M_A + M_B) + 4\sqrt{N(N+1)}\sqrt{M_A M_B} \cos(\gamma_A + \gamma_B - \varphi) - (M_A - M_B)^2}{2N(1 + M_A + M_B) + 4\sqrt{N(N+1)}\sqrt{M_A M_B} \cos(\gamma_A + \gamma_B - \varphi) + M_A + M_B}. \quad (24)$$

It shows a maximum non-classical violation of Lee criterion ($\mathcal{P}_{Lee}^{(C)} = 1$) when the intensities of seeds are null. Eventually, in the case of squeezed seeding one obtains

$$\mathcal{P}_{Lee}^{(S)} = \frac{2N(1 + N_A + N_B) - 2(N_A^2 + N_B^2) - (N_A - N_B)^2 - N_A - N_B}{2N(1 + N_A + N_B) + N_A + N_B}. \quad (25)$$

5 Entanglement

The downconversion process is known to provide pairwise entanglement between A - and B - beams. In the spontaneous process, as well as in the case of coherent and vacuum squeezed seeds the output state is entangled for any value of $N \neq 0$ whereas in the case of a thermally seeded PDC the degree of entanglement crucially depends on the intensity of the seeds, as shown in [31]. In fact, as ρ_{out} is a Gaussian state (since thermal states are Gaussian and the PDC Hamiltonian is bilinear in the field modes) its entanglement properties can be evaluated by checking the positivity of the partial transpose (PPT condition), which represents a sufficient and necessary condition for separability for Gaussian pairwise mode entanglement [35].

In order to check whether and when the state ρ_{out} is entangled we apply the PPT criteria for Gaussian entanglement [35]. For instance, we apply the positive map \mathcal{L}_B to the state ρ_{out} , $\mathcal{L}_B(\rho_{out})$ being the transposition (complex conjugation) only of the subspace B .

Gaussian states are completely characterized by their covariance matrix. In particular the covariance matrix of ρ_{out} is \mathbf{V} , with $V_{\alpha\beta} = 2^{-1}\langle\{\Delta w_\alpha, \Delta w_\beta\}\rangle$, where the vector operator $\mathbf{w} = (X_A, Y_A, X_B, Y_B)^T$, with the ‘‘position’’(-like) operators X and ‘‘momentum’’(-like) operators Y are $X_j = \frac{a_j + a_j^\dagger}{\sqrt{2}}$ $Y_j = \frac{a_j - a_j^\dagger}{i\sqrt{2}}$. Thus, the separability properties of ρ_{out} can be obtained from the positivity of $\mathcal{L}_B(\rho_{out})$, which can be expressed in terms of its covariance matrix $\tilde{\mathbf{V}}$ as

$$\tilde{\mathbf{V}} + \frac{i}{2}\mathbf{\Omega} \geq 0, \quad (26)$$

with

$$\mathbf{\Omega} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}. \quad (27)$$

Simon showed that $\tilde{\mathbf{V}}$ can be calculated exactly as \mathbf{V} with a sign change in the B momentum variable ($Y_B \rightarrow -Y_B$), while the other momentum and position variables remain unchanged [35]. Thus, we obtain

$$\tilde{\mathbf{V}} = \begin{pmatrix} \mathcal{A}_1 & \mathcal{D} & \mathcal{C}_1 & \mathcal{G}_1 \\ \mathcal{D} & \mathcal{A}_2 & \mathcal{G}_2 & \mathcal{C}_2 \\ \mathcal{C}_1 & \mathcal{G}_2 & \mathcal{B}_1 & \mathcal{F} \\ \mathcal{G}_1 & \mathcal{C}_2 & \mathcal{F} & \mathcal{B}_2 \end{pmatrix} \quad (28)$$

where in the thermal case

$$\begin{aligned}
\mathcal{A}_1 &= \mathcal{A}_2 = 1/2 + \mu_A + N(1 + \mu_A + \mu_B), \\
\mathcal{B}_1 &= \mathcal{B}_2 = 1/2 + \mu_B + N(1 + \mu_A + \mu_B), \\
\mathcal{C}_1 &= \mathcal{C}_2 = \sqrt{N(N+1)}(1 + \mu_A + \mu_B), \\
\mathcal{G}_1 &= \mathcal{G}_2 = \mathcal{D} = \mathcal{F} = 0.
\end{aligned} \tag{29}$$

In the case of laser seeds the results are the same as in the case of spontaneous PDC, i.e. the same of Eq. (29) with $\mu_A = \mu_B = 0$. In the case of squeezed seeds we have

$$\begin{aligned}
\mathcal{A}_1 &= 1/2 + N(2 + N_A + N_B) + \sqrt{N_A(1 + N_A)}(1 + N) + \sqrt{N_B(1 + N_B)}N \cos(2\Delta\varphi), \\
\mathcal{A}_2 &= 1/2 + N(2 + N_A + N_B) - \sqrt{N_A(1 + N_A)}(1 + N) - \sqrt{N_B(1 + N_B)}N \cos(2\Delta\varphi), \\
\mathcal{B}_1 &= 1/2 + N(2 + N_A + N_B) + \sqrt{N_B(1 + N_B)}(1 + N) + \sqrt{N_A(1 + N_A)}N \cos(2\Delta\varphi), \\
\mathcal{B}_2 &= 1/2 + N(2 + N_A + N_B) - \sqrt{N_B(1 + N_B)}(1 + N) - \sqrt{N_A(1 + N_A)}N \cos(2\Delta\varphi), \\
\mathcal{D} &= \sqrt{N_B(1 + N_B)}N \sin(2\Delta\varphi), \\
\mathcal{F} &= \sqrt{N_A(1 + N_A)}N \sin(2\Delta\varphi), \\
\mathcal{C}_1 &= \left[1 + N_A + N_B + \sqrt{N_A(1 + N_A)} + \sqrt{N_B(1 + N_B)} \right] \sqrt{N(1 + N)} \cos(\Delta\varphi), \\
\mathcal{C}_2 &= \left[-1 - N_A - N_B + \sqrt{N_A(1 + N_A)} + \sqrt{N_B(1 + N_B)} \right] \sqrt{N(1 + N)} \cos(\Delta\varphi), \\
\mathcal{G}_1 &= \left[1 + N_A + N_B + \sqrt{N_A(1 + N_A)} - \sqrt{N_B(1 + N_B)} \right] \sqrt{N(1 + N)} \cos(\Delta\varphi), \\
\mathcal{G}_2 &= \left[1 + N_A + N_B - \sqrt{N_A(1 + N_A)} + \sqrt{N_B(1 + N_B)} \right] \sqrt{N(1 + N)} \cos(\Delta\varphi),
\end{aligned} \tag{30}$$

with $\Delta\varphi = \zeta_A/2 + \zeta_B/2 - \varphi$.

The PPT criterion of Eq. (26) can be rewritten in terms of the smallest partially transposed symplectic eigenvalue \tilde{d}_- as $\tilde{d}_- \geq 2^{-1}$. This condition is never satisfied in the case of coherent and vacuum squeezed seeds, while in the case of thermal seeds we obtain that

$$\tilde{d}_- = \frac{1}{\sqrt{2}} \sqrt{\mathcal{A}_1^2 + \mathcal{B}_1^2 + 2\mathcal{C}_1^2 - \sqrt{(\mathcal{A}_1 + \mathcal{B}_1)^2[(\mathcal{A}_1 - \mathcal{B}_1)^2 + 4\mathcal{C}_1^2]}}, \tag{31}$$

thus the condition $\tilde{d}_- \geq 2^{-1}$ is satisfied when [31]

$$\mu_A \mu_B - N(1 + \mu_A + \mu_B) \geq 0. \tag{32}$$

It is noteworthy to observe that the separability/entanglement properties of the state ρ_{out} with thermal seeds can be highlighted by the direct photon counting on A- and B- arms. In fact, with an ideal detection system, the inequality

$$\langle [\Delta(n_A - n_B)]^2 \rangle - (\langle n_A \rangle - \langle n_B \rangle)^2 \leq \langle n_A \rangle + \langle n_B \rangle. \tag{33}$$

exactly corresponds to Eq. (32).

Thus, as in the two previous cases we can define a parameter quantifying the amount of the violation of the separability bound \mathcal{P}_{Ent} , only in the case of thermal seeds

$$\mathcal{P}_{Ent} = 1 - \frac{\langle [\Delta(n_A - n_B)]^2 \rangle - (\langle n_A \rangle - \langle n_B \rangle)^2}{\langle n_A \rangle + \langle n_B \rangle}. \quad (34)$$

$\mathcal{P}_{Ent} = 0$ corresponds to the boundary between the separability and the entanglement regions, in fact for the state ρ_{out} with thermal seeds we obtain

$$\mathcal{P}_{Ent} = 2 \frac{N(1 + \mu_A + \mu_B) - \mu_A \mu_B}{2N(1 + \mu_A + \mu_B) + \mu_A + \mu_B}. \quad (35)$$

According to Eq. (35) we observe that ρ_{out} is entangled when $0 < \mathcal{P}_{Ent} \leq 1$, and that the maximal violation of the separability bound (corresponding to $\mathcal{P}_{Ent} = 1$) is achieved by the spontaneous PDC ($\mu_A = \mu_B = 0$), while, if one of the two arms is seeded by the vacuum, irrespective of the magnitude of the thermal seed on the other arm, the state is always entangled.

We underline that the parameter \mathcal{P}_{Ent} cannot be considered an entanglement measure (as it does not have the correct properties) [36]. A quantification of entanglement which can be computed for general two-mode Gaussian states is provided by, e.g., the logarithmic negativity.

6 Discussion and conclusion

After having introduced the classicality quantifiers in the previous sections, here we compare them directly: fig. 1 shows the regions of nonclassicality of the three features considered, namely, the entanglement, the sub-shot noise, and the Lee's nonclassicality. In particular, in fig. 1 (a) the three regions are plotted in the case of thermal seeds, in fig. 1 (b) and (c) are in the case of coherent seeds with $\cos(\gamma_A + \gamma_B - \varphi) = 1$ and $\cos(\gamma_A + \gamma_B - \varphi) = -1$ respectively, and fig. 1 (d) in the case of squeezed vacuum seeds.

The most internal region corresponds to the ranges of values of seed mean number of photons (μ_A, μ_B for the thermal seeds, M_A, M_B for the coherent seeds, N_A, N_B for the squeezed vacuum seeds), and spontaneous PDC mean number of photons N where ρ_{out} is nonclassical in the context of Lee's criteria. It is noteworthy to observe that according to Eq. (9) and Eq. (18), being Lee-nonclassical is a sufficient condition to be sub-shot noise, as it can be seen also from Fig. 1. Furthermore, we note that in the case of coherent and vacuum squeezed seeds the output state is always entangled. In the thermal case, according to Eq. (9) and to Eq. (32), it can be observed that being sub-shot noise limited is a sufficient condition for being entangled (fig. 1 (a)). It is interesting to note that in the thermal case the three conditions coincide when $\mu_A = \mu_B$. Analogously, for the vacuum squeezed fields the SNL limit and the Lee's criterion limit converge when $N_A = N_B$, as in can be observed in fig. 1 (d). The same happens in the case of coherent seeds. In this case it is interesting to note that, as the output state always violate the SNL limit when $\cos(\gamma_A + \gamma_B - \varphi) \geq 0$, when $M_A = M_B$ also the Lee's criterion is always satisfied (see fig. 1 (b)).




Fig. 1 Regions of nonclassicality (entanglement, SNL violation, Lee's classicality criterion violation) plotted as a function of the mean number of photons of the seeding fields, and of the spontaneous PDC (N) for the different state of the seeding fields considered. In particular (a) thermal seeds, (b) coherent seeds with $\cos(\gamma_A + \gamma_B - \varphi) = 1$, (c) coherent seeds with $\cos(\gamma_A + \gamma_B - \varphi) = -1$, (d) vacuum squeezed seeds.

In conclusion, we have shown that there is a well defined hierarchy among the considered nonclassicality criteria (entanglement, SNL violation, Lee's classicality criterion violation) when PDC is seeded by simple single mode Gaussian states (such as thermal, coherent and vacuum squeezed states). The natural extension of this work is the investigation of such a hierarchy in the presence of the most general single mode Gaussian state as seeding fields. Work along these lines is in progress and results will be presented elsewhere [37].

References

1. T. Nieuwenhuizen et al. Editors, *Beyond The Quantum*, World Scientific, Singapore (2007); L. Accardi et al., *Foundations of Probability and Physics 5*, AIP, New York (2009).
2. M. Genovese, *Phys. Rep.* **413**, 319 (2005) and refs therein.
3. G. Auletta, "Foundations and Interpretation of Quantum Mechanics", (World Scientific, Singapore, 2000).
4. P. Ghose, *Testing Quantum Mechanics on a New Ground*, Cambridge Univ. Press, Cambridge (2000).
5. E. Santos, *Advanced Science Letters 2* (2009) 475 and other papers of the special issue.
6. M. Genovese, C. Novero and E. Predazzi, *Phys. Lett. B* **513**, 401 (2001).
7. S. Pirandola, *Phys. Rev. A* **68**, 0311186 (2003);
8. A. Serafini, F. Illuminati, M. G. A. Paris and S. De Siena, *Phys. Rev. A* **69**, 022318 (2004).
9. A. Serafini, M. G. A. Paris, F. Illuminati and S. De Siena, *J. Opt. B* **7**, R19 (2005).
10. A. Serafini, S. De Siena, F. Illuminati and M. G. A. Paris, *J. Opt. B* **6**, 591 (2004).
11. M. G. A. Paris, F. Illuminati, A. Serafini and S. De Siena, *Phys. Rev A* **68**, 012314 (2003)
12. G. M. D'Ariano, M. F. Sacchi and P. Kumar, *Phys. Rev. A* **59**, 826 (1999).
13. D. N. Klyshko, *Phys. Lett. A* **213**, 7 (1996).
14. Arvind, N. Mukunda and R. Simon, *J. Phys. A* **31**, 565 (1998).
15. W. Vogel, *Phys. Rev. Lett.* **84**, 1849 (2000).
16. Th. Richter, W. Vogel, *Phys. Rev. Lett.* **89**, 283601 (2002).
17. Th. Richter and W. Vogel, *J. Opt. B: Quantum Semiclass. Opt.* **5**, S371 (2003).
18. A. Lvovsky and S. Shapiro, *Phys. Rev. A* **65**, 033830 (2002).
19. B. J. Dalton, *Phys. Scr.* **T 12**, 43 (1986).
20. W. Schleich and J.A. Wheeler, *Nature (London)* **326**, 574 (1987).
21. S. Schiller et al., *Phys. Rev. Lett.* **77**, 2933 (1996).
22. V. Vedral and M.B. Plenio, *Phys. Rev. Lett.* **57**, 1619 (1998).
23. R. Alicki and N. Van Ryn, *J. Phys. A: Math. Theor.* **41**, 062001 (2008).
24. G. Brida, I. P. Degiovanni, M. Genovese, V. Schettini, S. Polyakov and A. Migdall, *Opt. Exp.* **16**, 11750 (2008); G. Brida, I. P. Degiovanni, M. Genovese, F. Piacentini, V. Schettini, N. Gisin, S. Polyakov and A. Migdall, *Phys. Rev. A* **79**, 044102 (2009).
25. T. Ishkakov et al., *JETP Lett.* **88**, 660 (2008).
26. N. Spagnolo et al., *Physical Review A* **80**, 032318 (2009)
27. A. Ourjuntsev et al., *Nature* **448**, 784 (2007).
28. G. Brida et al., arXiv:0904.3009; arXiv:0907.4117.
29. I. P. Degiovanni, M. Genovese, V. Schettini, M. Bondani, A. Andreoni and M. Paris, *Phys. Rev. A* **79**, 063836 (2009).
30. I. Ruo-Berchera, *Adv. Sci. Lett.*, **2**, 407 (2009).
31. I. P. Degiovanni, M. Bondani, E. Puddu, A. Andreoni and M. G. A. Paris, *Phys. Rev. A* **76**, 062309 (2007).

32. C. T. Lee, Phys. Rev. A **41**, 1569 (1990).
33. C. T. Lee, Phys. Rev. A **42**, 1608 (1990).
34. L. Mandel, Opt. Lett. **4**, 205 (1979).
35. R. Simon, Phys. Rev. Lett. **84**, 2726 (2000).
36. G. Adesso, F. Illuminati, J. Phys. A, **40**, 7821 (2007).
37. I. Degiovanni et al., *in preparation*.